



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>7</sup> :  A61K 31/555, 31/535, 31/495, 31/44, 31/40, 31/16, C07D 265/30, 241/00, 211/00, 207/00, C07C 231/00, 233/00		A1	(11) International Publication Number: <b>WO 00/32200</b>  (43) International Publication Date: <b>8 June 2000 (08.06.00)</b>
(21) International Application Number: <b>PCT/US99/28136</b>  (22) International Filing Date: <b>24 November 1999 (24.11.99)</b>		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(30) Priority Data:  60/109,615 24 November 1998 (24.11.98) US		Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(71)(72) Applicants and Inventors: MAKRIYANNIS, Alexandros [US/US]; 16 Stearns Road, Watertown, MA 02172 (US). KHANOLKAR, Atmaram [IN/US]; One South Eagleville Road #55, Storrs, CT 06269 (US). GOUTOPOULOS, Andreas [GR/US]; 146 Hunting Lodge, Storrs, CT 06269 (US).		(74) Agents: YALE, Guy, D. et al.; Alix, Yale & Ristas, LLP, 750 Main Street, Hartford, CT 06103 (US).	

(54) Title: CANNABIMIMETIC LIPID AMIDES AS USEFUL MEDICATIONS

## (57) Abstract

Novel analogs of arachidonylethanolamide are presented which have higher affinities for the cannabinoid CB1 and/or CB2 receptor sites. Further, most of the analogs exhibit greater metabolic stability than arachidonylethanolamide. The improved receptor affinity and selectivity and/or greater metabolic stability make these analogs therapeutically useful as medications for relief of pain caused by cancer and nausea caused by chemotherapy, as well as for peripheral pain. The compounds may also be useful as oral and topical contraceptives, in suppression of the immune system, enhancement of appetite and in treatment of psychomotor disorders, multiple sclerosis and hypertension.